

RESEARCH ARTICLE

PEER REVIEWED | OPEN ACCESS

Rehabilitation of a patient with spinal cord lesion due to surgical removal of the spinal tumor with chronic idiopathic demyelinating polyneuropathy (CIDP): A case report

Md Zakir Hossain, Akter Sohana, MD Waliul Islam, Shahid Afridi

ABSTRACT

Aims: The literature provides numerous medical therapies for chronic inflammatory demyelinating polyradiculoneuropathy (CIDP) and spinal cord lesion following spinal tumor surgery separately. Spinal cord injuries from CIDP-related spinal tumors have no evidence-based rehabilitation methods. This case report discusses CIDP, spinal cord lesion owing to a spine tumor, and tertiary care interdisciplinary rehabilitation. It shows how medical knowledge, clinical reasoning, and evidence guide outcome measures, care plans, and clinical decisions help to overcome CIDP.

Methods: In this report, we follow a 27-year-old male who began experiencing gradual paresthesia and mild weakening in his lower extremities for six weeks. After two months, the lumbar spine had the initial operation (laminectomy) because of the extreme pain. After the operation, he was pain-free, able to walk, and even started riding again. Pain in his lower midback, similar to cramping, and moderate swelling in his left ankle forced him into a wheelchair three years after his initial operation (intradural-extramedullary spinal space-occupying lesion at L2–L5 level). After the second operation (laminectomy), he had trouble in walking, lost sensation below the knees on both legs, and experienced mild incontinence. As the patient was confined to a

wheelchair, he was standing with the help of two people. Significant main muscle group weakening was one of the first noticeable symptoms. Therapeutic exercise, balance training, functional training, and progressive endurance activities were the main components of the intervention. Berg balance score increased from 5 to 23, which is an eligible score for discharge.

Results: Clinical outcomes for the man with spinal cord lesion and chronic inflammatory demyelinating polyradiculoneuropathy (CIDP) were improved with interdisciplinary therapy. The patient underwent a first lumbar spine operation, which initially eased pain and allowed him to walk and ride again after feeling gradual paresthesia and minor weakened in his lower extremities.

Conclusion: For this patient with CIDP with spinal cord lesion due to spinal tumor, effective collaborative team communication and interdisciplinary management worked to optimize clinical decision making and recovery.

Keywords: Chronic inflammatory demyelinating polyneuropathy, Demyelinating conditions, Rehabilitation, Spinal tumor

How to cite this article

Hossain MZ, Sohana A, Islam MW, Afridi S. Rehabilitation of a patient with spinal cord lesion due to surgical removal of the spinal tumor with chronic idiopathic demyelinating polyneuropathy (CIDP): A case report. Edorium J Disabil Rehabil 2023;9(2):15–22.

Article ID: 100056D05MH2023

doi: 10.5348/100056D05MH2023RA

Md Zakir Hossain¹, Akter Sohana², MD Waliul Islam¹, Shahid Afridi¹

Affiliations: ¹Centre for the Rehabilitation of the Paralyzed (CRP), Savar, Dhaka, Bangladesh; ²Clinical Physiotherapist, Centre for the Rehabilitation of the Paralyzed, Dhaka, Bangladesh.

Corresponding Author: Md Zakir Hossain, Centre for the Rehabilitation of the Paralyzed (CRP), Savar, Dhaka, Bangladesh; Email: physio.zakir@gmail.com

Received: 14 July 2023

Accepted: 02 October 2023

Published: 17 November 2023

INTRODUCTION

Intradural or extradural spinal tumors originate from the spinal cord or vertebrae. Intradural tumors are intramedullary, extramedullary, or both [1–3]. Most primary spinal cord cancers are schwannomas, generated from Schwann cells. Slow-growing, benign tumors can compress the spinal cord and nerve roots. Schwannomas are more common in the lumbar, thoracic, and cervical spine, and frequently detected after the age of 30 years. Chronic idiopathic demyelinating polyneuropathy (CIDP), an immune-mediated neuropathy, causes increasing weakening, sensory impairment, and missing or decreased tendon reflexes [4]. It occurs 0.67–10.3 times per 100,000 people [5]. Chronic idiopathic demyelinating polyneuropathy can be asymmetric, focal, or sensory/motor-dominated. European Federation of Neurological Societies/Peripheral Nerve Society 2010 CID diagnostic criteria are extensively utilized [6,7]. Chronic idiopathic demyelinating polyneuropathy treatment tries to minimize inflammation, demyelination, and secondary axonal degeneration. Therapies continue until stabilization or maximum improvement [8]. Sensation, strength, and daily life gains indicate treatment efficacy. Infections, systemic disorders, and neurotoxic medications might worsen CIDP symptoms [9]. A 12-week aerobic or resistance exercises program for CIDP patients. The training program targeted specific muscle regions, whereas the aerobic program comprised regular cycling. Both exercise programs improved function and reduced symptoms [10]. Trauma, spinal degeneration, tumor compression, vascular disease, infection, and inflammation can cause spinal cord injury (SCI). Spinal cord injury treatment aims to improve symptoms, function, and long-term remission. Using the International Classification of Functioning, Disability, and Health (ICF) framework, physiotherapists can develop effective rehabilitation strategies for patients with neurological conditions by understanding the characteristics and management of spinal cord lesions with CIDP.

MATERIALS AND METHODS

The case has been selected conveniently in neurology unit of Centre for the Rehabilitation of the Paralyzed (CRP). The duration of observation was six months. The cases have been presented partially followed by case report (CARE) guideline and verbal consent has been taken from the patients and caregiver of the patients.

Case summary

Mr. Wahid, then 27 years old, began experiencing paresthesia and slight weakness in his lower limbs in 2013. He had suffered from lower back swelling ever since he was a kid, but he had managed to keep working through it. In 2013, however, he developed significant low back

pain that rendered him bedridden; this ultimately led to lumbar spine surgery. The operation was fruitful, and he is now pain-free, mobile, and able to resume walking and cycling. The pain in Mr. Wahid's lower midback and some moderate edema in his left ankle in 2016 rendered him wheelchair-bound once more. As a result, a second operation was required to remove an intradural-extramedullary spinal space-occupying lesion between lumbar levels L2 and L5. He developed slight bowel and bladder incontinence and was unable to walk following the second operation. Two people helped him to get up on his feet. Mr. Wahid's symptoms worsened during the next six months, including upper- and lower-limb paresthesia and sleep jerking. Mr. Wahid had a hard time finding appropriate treatment for his symptoms despite the fact that he had no known medical disorders such as diabetes, hypertension, thyroid issues, kidney disease, or heart disease. Following visits to the Neuroscience Hospital in Bangladesh and the Christian Medical College in India, among others, he was referred to physiotherapy specialists for further care and rehabilitation. In 2018, Mr. Wahid's condition started to improve after 15 days of physiotherapy at Mymensingh Medical College Hospital. His rehabilitation showed promise and provided hope for more progress in the future, despite his ongoing reduced lower limb sensation, weakness in both legs, poor standing balance, urinary concerns, coordination challenges, and difficulty managing his stride.

Assessment

International Classification of Functioning, Disability, and Health (ICF) requires understanding disability state to improve a person's quality of life [11]. Managing a person with any type of disability is multi-dimensional and comprehensive. The patient complained of significant weakness in both lower limbs and moderate weakness in the upper limb, modest upper limb pain, inability to stand, bladder problems, walking, and difficulties with activities of daily living (ADLs). His elbow pain, severe pelvic control, trunk control, wrist, and lower limb weakness with foot drop and moderate muscular wasting in tibialis anterior and surrounding the foot, gait irregularity, and upper limb fine motor activity were assessed. Visual analogue scale (VAS) pain was 6 out of 10 and decreased lower limb active range of motion (Hip Flexion 20°, abduction loss, 50° knee flexion present with knee extension full, ankle, foot movement in both lower limbs absent, upper limb range of motion is full). Lower limbs are withering. After two years, he went to physiotherapy. American Spinal Injury Association (ASIA) exam was his initial evaluation.

Instrumentation and tools

The American Spinal Injury Association assessed the case. This instrument is valid and reliable for measuring

spinal cord injury neurological level and outcome [12]. CRP physiotherapy department neurology unit self-assessment is also included. Personal details included demographic information, history of present complaint, patient complaint, past medical history, medications, investigation, social history, home environment, general observations like consciousness, vision, hearing, edema, drooling, speech, subluxation, local observations like blood pressure, temperature, respiratory function, posture, upper and lower limb function uncompressing tonicity, sensation, and proprioception.

RESULTS

Clinical outcomes for the man with spinal cord lesion and chronic inflammatory demyelinating polyradiculoneuropathy (CIDP) were improved with interdisciplinary therapy. The patient underwent a first lumbar spine operation, which initially eased pain and

allowed him to walk and ride again after feeling gradual paresthesia and minor weakened in his lower extremities. But by the age of 3 years, he was confined to a wheelchair due to lower midback pain and considerable swelling in his left ankle (Table 1). After undergoing a second procedure, the patient experienced slight incontinence, difficulty walking, and a loss of sensation below the knees (Table 2). Therapeutic exercise, balancing instruction, functional drills, and progressive endurance activities made up the rehabilitation intervention (Table 3).

The patient's Berg balance score improved greatly from 5 to 23, making him a candidate for discharge (Table 4). When the interdisciplinary team worked together and communicated well, it helped guide clinical decision-making and speed the patient's recovery (Table 5). This case report serves as a useful resource for future instances of a similar nature, highlighting the significance of evidence-based rehabilitation strategies in the management of spinal cord injury caused by CIDP-related spine tumors (Figure 1).

Table 1: Different types of spinal tumors

| Vertebral column tumors (extradural) | Spinal cord tumors (intradural) |
|--------------------------------------|-----------------------------------|
| Primary | Primary |
| Malignant tumors | Intramedullary tumors |
| Osteosarcoma | Astrocytoma |
| Chondrosarcoma | Ependymoma |
| Fibrosarcoma | Dermoid tumor |
| Malignant fibrosis histiocytoma | Epidermoid tumor |
| Ewing's sarcoma | Teratoma |
| Multiple myelomas | Lipoma |
| Lymphoma | Hemangioblastoma |
| Chordoma | Ganglioglioma |
| | Oligodendroglioma |
| Benign tumors | Extramedullary tumors |
| Osteoid osteoma | Meningioma |
| Osteoblastoma | Neurofibroma (nerve sheath tumor) |
| Osteochondroma | Schwannoma (nerve sheath tumor) |
| Enchondroma | Dumbbell tumors |
| Chondroblastoma | Secondary |
| Chondromyxoid fibroma | Metastatic tumors |
| Fibroma | |
| Giant cell tumor Hemangioma | |
| Aneurysmal bone cyst | |
| Eosinophilic granuloma | |
| Secondary | |
| Metastatic tumors | |

Table 2: Initial ASIA assessment and after six months of treatment

| | | |
|------------------------------------|-----------------------|-------------------|
| Skeletal level | | L2–L5 |
| Neurological levels | Sensory right side L3 | |
| Neurological levels | Sensory left side L3 | |
| Neurological levels | Motor right side L3 | |
| Neurological levels | Motor left side L3 | |
| Neurological level of injury (NLI) | | L3 |
| Complete/Incomplete | | Sensory preserved |
| ASIA diagnosis | Incomplete | ASIA B |
| Outcome after six months | | |
| Skeletal level | | L2–L5 |
| Neurological levels | Sensory right side L3 | |
| Neurological levels | Sensory left side L3 | |
| Neurological levels | Motor right side L3 | |
| Neurological levels | Motor light side L3 | |
| Neurological level of injury (NLI) | | L3 |

Table 3: Impairments of cases in initial assessment

| Body structure and functions | Activity limitations | Participation restrictions | Personal factors | Environmental factors |
|---|---|--|--|--|
| <ul style="list-style-type: none"> Decrease active ROM Foot drop Muscle strength: <ul style="list-style-type: none"> Upper limb = 4 oxford muscle grade Rt LL-table 4 Lt LL-table 4 Sensory intact up to L3 Bowel bladder impairment (sensory present, motor problem) | <ul style="list-style-type: none"> Wheelchair bounded Poor sitting on a bed Unable to stand without maximal support Unable to walk Unable to control urination and micturition Unable to feel in lower limb | <ul style="list-style-type: none"> Drooped from job Restrictions in engaging with the friend circle. The attitudinal problem of relatives due to lack of education. | <ul style="list-style-type: none"> The elder son in the family Young age Depression | <ul style="list-style-type: none"> Lives in rural area Environmental inaccessibility |

Table 4: Manual muscle testing of major muscle group

| Muscle groups | Initial examination | Discharge |
|----------------------|---------------------|-----------|
| Shoulder flexion | 3/5 | 4/5 |
| Shoulder abduction | 3/5 | 4/5 |
| Elbow flexion | 3/5 | 5/5 |
| Elbow extension | 3/5 | 5/5 |
| Wrist flexion | 3/5 | 5/5 |
| Wrist extension | 3/5 | 5/5 |
| Finger flexion | 2/5 | 4/5 |
| Finger abduction | 2/5 | 4/5 |
| Hip flexion | 2/5 | 4/5 |
| Hip extension | 1/5 | 3/5 |
| Knee flexion | 0/5 | 1/5 |
| Knee extension | 0/5 | 3/5 |
| Ankle dorsiflexion | 0/5 | 0/5 |
| Ankle plantarflexion | 0/5 | 0/5 |

Table 5: Summary of physical therapy intervention, AAROM: active-assisted range of motion; PROM: passive range of motion; AROM: active range of motion; ROM: range of motion; AFO: ankle-foot orthosis

| Month 1 | Month 2 | Month 3 | Month 4 | Month 5 | Month 6 |
|--|--|---|---|---|---|
| <ul style="list-style-type: none"> Positioning, passive stretching, facilitating sensory integration practice to improve tone PROM and active-assisted ROM to upper and lower extremities, [10] repetitions each Bridging exercise, Sitting exercise and functional activities Standing frame Kegel exercise | <ul style="list-style-type: none"> AAROM and AROM to upper and lower extremities, 5-10 repetitions each Sitting balance and functional activities Standing Frame Transfer training between bed and wheelchair Motor training exercise in limbs, pelvis, trunk in lying, sitting, standing [13] Bridging exercise Kegel exercise | <ul style="list-style-type: none"> AAROM and AROM to upper and lower extremities, 5-10 repetitions each Sitting balance and functional activities Standing Frame Transfer training between bed and wheelchair A motor training exercise in limbs, pelvis, trunk in lying, sitting, standing [13] Kegel exercise | <ul style="list-style-type: none"> Trunk and pelvic control exercise in lying progress to four points lying, half-lying, sitting and supported standing [13]. Bridging exercise Standing balance by balance board and functional activities at the edge of the bed Kegel exercise | <ul style="list-style-type: none"> Balance exercise in four-point kneeling, sitting and standing [14] Strength training in lower limbs, abdomen, and pelvis [14] Gait training from the maximally supported four-point walker, parallel bar, and progressed to elbow crutch [14] Solid AFO for improving gait | <ul style="list-style-type: none"> Balance exercise in four-point kneeling, sitting and standing [15] Strength training in lower limbs, abdomen, and pelvis [16] Gait training from the maximally supported four-point walker, parallel bar, and progressed to elbow crutch with AFO [16]. |



Figure 1: Patient standing in parallel bar without support.

DISCUSSION

As each condition affects symptom management, treatment, rehabilitation, and prognosis, having a spinal cord lesion (SCL) from tumor removal and chronic inflammatory polyneuropathy (CIDP) can be devastating for patients and their families. In a study, spinal cord tumors are abnormal growths of tissue that can compress or invade the spinal cord or its surrounding structures, causing neurological deficits, pain, and disability and surgical removal of the tumor is often the primary treatment option, but it may also cause

damage to the spinal cord or its blood supply, resulting in further impairment. The patient made minimal physical development in the first month of physiotherapy. His initial pain, low function, weak endurance, and mobility anxiety hampered physical therapy approaches. In a study, CIDP is a rare autoimmune disorder that causes progressive inflammation and demyelination of the peripheral nerves, leading to weakness, numbness, and sensory loss. Chronic idiopathic demyelinating polyneuropathy can coexist with spinal cord tumors, either as a paraneoplastic syndrome or as an incidental finding. Chronic idiopathic demyelinating polyneuropathy may also worsen the neurological outcome after spinal cord tumor surgery. The physical therapist's supportive treatment, persistence, encouragement, and reinforcement of even tiny advances likely reduced functional deterioration and immobility, laying the groundwork for recovery. The physical therapist monitored the patient's condition and implemented a flexible care plan using knowledge of CIDP and spinal cord damage and literature. Physical therapy is geared toward muscle strengthening, communication, and mobility. It may involve exercises, stretching, massage, electrical stimulation, hydrotherapy, and gait training. Medical therapies reversed the inflammatory process, allowing recuperation. Despite strength and endurance gains, the patient showed slow functional progress.

CONCLUSION

The development of standardized outcome measures and specific practice guidelines for CIDP with spinal cord lesion could lead to best-practice care for this condition.

REFERENCES

1. Beall DP, Googe DJ, Mitchell BS. Intramedullary spinal cord tumors. *The Surgical Clinics of North America* 2007;87(5):1117–33.
2. Chongyi F, Yonggang W, Xiaoguang L, Zhongjun L, Junming M, Qingsan Z. Analysis of the clinical symptoms and surgical treatment of 95 cases of intradural-extramedullary tumors. *Journal of Clinical Neuroscience* 2008;15(5):495–8.
3. Arnaudovic KI, Gokaslan ZL. Surgery for spinal tumors: Indications and techniques. *Journal of Neuro-Oncology* 2019;143(3):387–94.
4. Elia-Pasquet S, Franques J, Devaux J, Fauchais AL, Blaison G. Chronic inflammatory demyelinating polyneuropathy. *Revue Neurologique* 2004;160(12 Pt 1):1223–30.
5. Schellinger KA, Propp JM, Villano JL, McCarthy BJ. Descriptive epidemiology of primary spinal cord tumors. *J Neurooncol* 2008;87(2):173–9.
6. Sowash MG, Ballinger WE, Juel VC. Chronic inflammatory demyelinating polyneuropathy: An update on diagnosis and management. *Clinical Neurology and Neurosurgery* 2017;154:98–106.
7. Milano E, Bianco M, Amoroso A, Di Sapio A, Sessa E, Toscano A. Chronic inflammatory demyelinating polyradiculoneuropathy: Diagnosis and treatment. *Neurological Sciences* 2010;31(5):573–81.
8. Broers MC, Bunschoten C, Nieboer D, Lingsma HF, Jacobs BC. Guillain-Barré syndrome subtype diagnosis: A validation study. *Journal of Neurology* 2019;266(12):3064–70.
9. Van den Bergh PY, Hadden RD, Bouche P, Cornblath DR. Peripheral nervous system involvement in immune-mediated diseases. *Handbook of Clinical Neurology*. Elsevier; 2010, Vol. 97, p. 663–80.
10. Markvardsen LH, Overgaard K, Heje K, Sindrup SH, Christiansen I, Jensen TS. Aerobic training in patients with chronic inflammatory neuropathies: A randomized controlled trial. *Brain and behavior* 2017;7(3):e00649.
11. Childs J, Blyth C, Pelletier C. *Physiotherapy Management of Lower Limb Conditions: International Classification of Functioning, Disability, and Health (ICF)*. London: Elsevier Health Sciences; 2008.
12. Marino RJ, Graves DE. Metric properties of the ASIA motor score: Subscales improve correlation with functional activities. *Arch Phys Med Rehabil* 2004;85(11):1804–10.
13. Ginis KM, Hicks AL, Latimer AE, et al. The development and testing of an exercise DVD for individuals with spinal cord injury. *Spinal Cord* 2018;56(9):836–43.
14. Tse MA, Tang A, Baumgartner L, Weir PL, Inness EL, Jaglal SB. Physical activity programming in the rehabilitation of persons with spinal cord injury: Defining strategies for success. *Spinal Cord* 2018;56(3):225–33.
15. Bye R, Pushman H, Summers J, et al. High-intensity resistance and functional training improve muscle strength, function, and disability in individuals with chronic incomplete spinal cord injury: A randomized controlled trial. *J Neurotrauma* 2017;34(1):23–9.
16. Mehrholz J, Kugler J, Pohl M. Locomotor training for walking after spinal cord injury. *Cochrane Database Syst Rev* 2012;11:CD006676.

Acknowledgments

The authors thank Md. KM Arman Hossain, Jashore University of Science & Technology for his valuable suggestion and guidelines and also extend their heartfelt thanks to Md. Akhlasur Rahman, Junior consultant, Spinal Cord Injury Unit, CRP for his technical support.

Author Contributions

Md Zakir Hossain – Conception of the work, Design of the work, Acquisition of data, Analysis of data, Interpretation of data, Drafting the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Akter Sohana – Conception of the work, Acquisition of data, Interpretation of data, Drafting the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

MD Waliul Islam – Conception of the work, Acquisition of data, Analysis of data, Drafting the work, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Shahid Afridi – Conception of the work, Design of the work, Acquisition of data, Analysis of data, Interpretation of data, Drafting the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Guarantor of Submission

The corresponding author is the guarantor of submission.

Source of Support

None.

Consent Statement

Written informed consent was obtained from the patient for publication of this article.

Conflict of Interest

Authors declare no conflict of interest.

Data Availability

All relevant data are within the paper and its Supporting Information files.

Copyright

© 2023 Md Zakir Hossain et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

ABOUT THE AUTHORS

Article citation: Hossain MZ, Sohana A, Islam MW, Afridi S. Rehabilitation of a patient with spinal cord lesion due to surgical removal of the spinal tumor with chronic idiopathic demyelinating polyneuropathy (CIDP): A case report. Edorium J Disabil Rehabil 2023;9(2):15–22.



Zakir Hossain is a Senior Clinical Physiotherapist at Neurology Unit, Centre for the Rehabilitation of the Paralyzed (CRP). He earned the undergraduate degree and Masters in Physiotherapy (MPT) from Bangladesh Health Professions Institute–BHPI (Academic Institute of CRP) from Physiotherapy Department. He has completed Bobath advanced course on Neurological condition. He has published several research papers in international academic journals. His research interests include rehabilitation and physiotherapy.

Email: Physio.zakir@gmail.com



Sohana Akter is a Clinical Physiotherapist at Neurology Unit, Centre for the Rehabilitation of the Paralyzed (CRP) for 5 years. She graduated and is continuing Masters in Physiotherapy (MPT Part-II) from the Physiotherapy Department of Bangladesh Health Professions Institute–BHPI (Academic Institute of CRP). She has finished the Advanced Bobath course on Neurological condition. She is still conducting post-graduation research on stroke rehabilitation a randomized clinical trial.

Email: sohana.mehejabin@gmail.com



MD Waliul Islam is a Senior Clinical Physiotherapist at Neurology Unit, Centre for the Rehabilitation of the Paralyzed (CRP). He earned the undergraduate degree and Masters in Physiotherapy (MPT) from Bangladesh Health Professions Institute–BHPI (Academic Institute of CRP) from Physiotherapy Department. He has published several research papers in international academic journals. His research interests include rehabilitation and physiotherapy.

Email: olee32@gmail.com



Shahid Afridi is a Clinical Physiotherapist at Centre for the Rehabilitation of the Paralyzed (CRP). He earned the undergraduate degree from Bangladesh Health Professions Institute–BHPI (Academic Institute of CRP) from Physiotherapy Department. He has published several research papers in international academic journals. His research interests include rehabilitation and physiotherapy.

Email: afridisbd2@gmail.com

Access full text article on
other devices



Access PDF of article on
other devices





INTERNATIONAL JOURNAL OF
CASE REPORTS AND IMAGES



VIDEO JOURNAL OF
CLINICAL RESEARCH



VIDEO JOURNAL OF
BIOMEDICAL SCIENCE




INTERNATIONAL JOURNAL OF
HEPATOBIILIARY AND
PANCREATIC DISEASES



INTERNATIONAL JOURNAL OF
BLOOD TRANSFUSION AND
IMMUNOHEMATOLOGY



EDORIUM JOURNAL OF
OPHTHALMOLOGY



EDORIUM JOURNAL OF
MEDICINE



EDORIUM JOURNAL OF
CARDIOTHORACIC AND
VASCULAR SURGERY



JOURNAL OF CASE REPORTS
AND IMAGES IN ORTHOPEDICS
AND RHEUMATOLOGY



EDORIUM JOURNAL OF
PSYCHOLOGY



EDORIUM JOURNAL OF
CELL BIOLOGY



JOURNAL OF CASE REPORTS AND IMAGES IN
DENTISTRY



EDORIUM JOURNAL OF
CANCER



EDORIUM JOURNAL OF
PSYCHIATRY



JOURNAL OF CASE REPORTS AND
IMAGES IN INFECTIOUS DISEASES



EDORIUM JOURNAL OF
ANATOMY AND EMBRYOLOGY



EDORIUM JOURNAL OF
SURGERY



JOURNAL OF CASE REPORTS
AND IMAGES IN PATHOLOGY



EDORIUM JOURNAL OF
ANESTHESIA